



Hopkins, J. C. A., Blazeby, J. M., Rogers, C. A., & Welbourn, R. (2016). The use of adjustable gastric bands for management of severe and complex obesity. *British Medical Bulletin*, 118(1), 67-75. <https://doi.org/10.1093/bmb/ldw012>

Peer reviewed version

Link to published version (if available):  
[10.1093/bmb/ldw012](https://doi.org/10.1093/bmb/ldw012)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via OUP at <https://academic.oup.com/bmb/article/118/1/64/1744476#35724673>. Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**The use of adjustable gastric bands for management of severe and complex obesity**

Journal:	<i>British Medical Bulletin</i>
Manuscript ID	BMB-2015-054.R1
Manuscript Type:	Invited Review
Date Submitted by the Author:	08-Feb-2016
Complete List of Authors:	Welbourn, Richard ; Musgrove Park Hospital, Hopkins, J; Fellow in Upper GI and Bariatric Surgery Blazeby, J Rogers, C
Keywords:	Obesity surgery, Bariatric surgery, Metabolic surgery, Gastric banding

SCHOLARONE™  
Manuscripts

Review

**The use of adjustable gastric bands for management of severe and complex obesity**

James C A Hopkins DPhil FRCS<sup>1</sup>,

Jane M Blazeby MD FRCS <sup>2</sup>

Chris A Rogers PhD <sup>3</sup>

Richard Welbourn MD FRCS<sup>1</sup>

<sup>1</sup> Department of Upper Gastrointestinal Surgery and Bariatric Surgery, Musgrove Park Hospital, Taunton TA1 5DA, UK

<sup>2</sup> Centre for Surgical Research, School of Social and Community Medicine, University of Bristol, Bristol

<sup>3</sup> Clinical Trials and Evaluation Unit, School of Clinical Sciences, University of Bristol, Bristol, UK

**Number of Figures = 1**

**Number of Colour Figures = 1**

**Number of Tables = 0**

Figure 1 has been previously published and permission has been given from Elsevier.

There is no supplementary material to go online.

**Running title:** Adjustable gastric banding for severe and complex obesity

**Keywords:** Obesity surgery, bariatric surgery, diabetes treatment

**Author for correspondence:** Mr Richard Welbourn, Consultant Surgeon, Dept Upper GI and Bariatric Surgery, Musgrove Park Hospital, Taunton TA1 5DA, UK

Richard.Welbourn@tst.nhs.uk, Tel +44 (0)1823 342100, Fax +44 (0)1823 343560

**Other authors' details:**

JCA Hopkins, Fellow in Upper GI and Bariatric Surgery, frog@doctors.org.uk

JM Blazeby, Professor of Surgery, J.M.blazeby@bristol.ac.uk

CA Rogers, Reader in Medical Statistics, Chris.Rogers@bristol.ac.uk

RW acts as guarantor and confirms that the manuscript has been submitted solely to this journal and is not published, in press, or submitted elsewhere.

**Conflict of Interest declaration:** James Hopkins, Jane Blazeby and Chris Rogers declare no conflict of interest. Richard Welbourn received sponsorship for attending conference/educational workshops from Allergan and Ethicon Endo-Surgery and is a consultant for Novo Nordisk. Musgrove Park Hospital, Taunton UK, also received grant funding for a bariatric surgery training fellowship from Ethicon Endo-Surgery.

Key words: obesity surgery, bariatric surgery, metabolic surgery, gastric banding

**ABSTRACT**

**Background**

Obesity levels in the UK have reached a sustained high and about 5% of the population would be candidates for bariatric surgery based upon current UK NICE guidelines, which has important implications for Clinical Commissioning Groups.

**Sources of data**

Summary data from Cochrane systematic reviews, randomised controlled trials (RCTs) and cohort studies.

**Areas of agreement**

Currently the only treatment that offers significant and durable weight-loss for those with severe and complex obesity is surgery. Three operations account for 95% of all bariatric surgery in the UK, but the NHS offers surgery to only a small fraction of those who could benefit. Laparoscopic adjustable gastric banding (gastric banding) has potentially the lowest risk and up-front costs of the three procedures.

**Areas of controversy**

Reliable level 1 evidence of the relative effectiveness of the operations is lacking.

**Growing points**

As a point intervention, weight-loss surgery together with the chronic disease management strategy for obesity can prevent significant future disease and mortality, and the NHS should embrace both.

**Areas timely for developing research**

Better RCT evidence is needed including clinical effectiveness and economic analysis to answer the important question 'which is the best of the three operations most frequently

performed?' This review considers the current evidence for gastric banding for the treatment of severe and complex obesity.

For Peer Review

INTRODUCTION

Obesity is a major worldwide public health problem with significant implications for primary care, particularly where obesity prevalence in developed countries is as high as 32.5% in women in high income North American countries, 29.8% in Australasia and 21.0% in Western Europe [1]. Although obesity levels may have reached a plateau, the burden of the important obesity related co-morbidities, such as type 2 diabetes, which was 382 million people worldwide in 2013, is predicted to rise to over 590 million in 2035 [2]. Both diabetes and obesity are independent risk factors for many important cancers [3, 4]. Obesity significantly increases many functional disorders and worsens overall health related quality of life [5], which can be improved by weight loss and metabolic surgery [6]. Overall, there is substantial evidence that obesity is a chronic disease that shortens life expectancy, even with the current best medical management of its co-morbidities [7].

Bariatric surgery is the only reproducibly effective treatment for severe and complex obesity [8, 9]. UK NICE guidelines provide clear guidance for the delivery of health-care and recommend that: bariatric surgery is the first line of treatment for those with a BMI of  $\geq 50$  in whom lifestyle interventions have failed; and those with a BMI  $\geq 40$ , or  $\geq 35$  with severe obesity-related co-morbidities, should be offered bariatric surgery [10]. Weight-loss and metabolic surgery procedures are among the most commonly performed elective general surgical operations in the world, with over 350,000 performed in 2011, [11] but in 2013 in the UK, with government funded health care, only around 8,000 operations were performed [12, 13], of which around 4,300 operations are recorded as being performed in the public sector [14].

Bariatric surgery is a young surgical speciality, with increasing numbers of cases being performed over a relatively short time frame [11], and there is a lack of high quality long term outcome data for the procedures. The most up-to-date, 2014, Cochrane collaboration systematic review of weight loss surgery in adults could only provide synthesizable evidence

for 3 years of follow up, and commented that ‘the long-term effects of surgery remain unclear’ [8]. However this level 1 evidence for short and medium term results of surgery can be compared directly to results for non-surgical interventions, and combined with some long-term cohort studies of bariatric surgery, albeit in smaller numbers, to show a pattern of long-term benefits from surgery. Given the established long-term benefit it is important to have large scale, pragmatic randomised controlled surgical trials to evaluate different techniques, to include economic evaluations, to give us the answer as to what is the best operation [15].


### **Gastric banding and how it facilitates weight loss**

In the UK >99% of gastric band procedures are performed laparoscopically and patients typically spend ≤1 days in hospital with very few early complications [12, 13]. The operation induces and sustains weight loss by activating a satiety mechanism, by adding or removing fluid from the band to achieve optimal filling. Food is not retained above the optimally adjusted band, i.e. not physically restricted, but briefly delayed food bolus transit through the band and the continuous pressure of the optimally filled band on the stomach wall produces both early satiety and a lack of appetite [16]. The effects of the band on oesophageal and proximal gastric function appears to activate a satiety signal, transmitted to CNS satiety centres via the vagus nerve, without physically restricting the meal size [16]. Figure 1 shows the position of a gastric band with a small “virtual” pouch of stomach below the gastro-oesophageal junction, above the final band position, held within gastro-gastro tunnelling sutures. Ideally patients are followed up every 1 – 3 months for the first two years by the bariatric multidisciplinary team (MDT), and then yearly in conjunction with GPs as part of a shared care model of chronic disease management (NICE CG189) [10].

Patients require life-long follow up for optimal weight control and co-morbidity resolution for gastric banding to work, especially as nutritional issues or complications can occur at any point after surgery [17, 18]. Symptoms of reflux, regurgitation or vomiting and especially nocturnal aspiration or dysphagia should alert the primary care GP to possible



complications. Abdominal pain along with any of these symptoms may suggest an acute complication, but other causes of abdominal pain should always be considered. Lifelong care involves essential counselling about food choices and eating patterns, as well as adjustment of the band [18]. Patients should therefore not be offered revisional surgery unless a process for continuing care is in place with the bariatric multidisciplinary team and within primary care.



**Figure 1** The adjustable gastric band. (Images reproduced from SM Griffin, SA Raimes, J Shenfine. Oesophagogastric Surgery. 5th ed. Saunders Elsevier, 2013)

**Weight loss effects of gastric bands & bariatric surgery**

The 2014 Cochrane review into weight loss surgery for adults included 22 randomised trials with 1,798 participants. The authors compared surgery with non-surgical interventions or different surgical procedures and most studies followed participants for only 12 to 36 months [8]. All of the trials which compared surgery with non-surgical interventions for weight loss found that surgery was significantly superior for weight loss at 1-2 years. This mirrored a 2013 systematic review and meta-analysis that directly compared surgery and medical therapy, using 11 randomised studies with 796 included patients, and a random effect model, clearly favouring surgery [9]. With study follow up of up to 2 years individuals

undergoing bariatric surgery lost more body weight (mean difference -26 kg (95% confidence interval -31 to -21),  $P < 0.001$ ) than those with medical therapy alone.

In the Cochrane review no clinically significant differences between the three main techniques of gastric band, all types of gastric bypass or sleeve gastrectomy existed for the majority of time points within 'multi-trial' evidence. Although only three trials directly compared gastric banding with laparoscopic Roux-en-Y gastric bypass (gastric bypass), the latter produced greater weight loss, with a mean difference in body mass index (BMI) reduction up to five years of -5.2 kg/m<sup>2</sup> (95% confidence interval -6.4 to -4.0;  $P < 0.00001$ ). However, this was only from a total of 265 patients when combined, and the quality of the evidence was described as *moderate* at best. In the seven included trials of gastric bypass and sleeve gastrectomy there was no consistent picture as to which procedure was better or worse, but with a *low* quality of evidence [8]. For the 2013 systematic review the cumulative body weight loss was similar but not statistically significantly different between gastric banding and gastric bypass or other techniques combined (mean difference between 6 and 7 kg,  $P > 0.20$ ) [9]. Effects on co-morbidities, complications and additional surgical procedures were neutral for the three main procedures in these reviews [8, 9]. Similarly a large prospective database, the Michigan Bariatric Surgery Collaborative, concluded that there was comparable effectiveness of gastric banding, gastric bypass and sleeve gastrectomy for the treatment of severe obesity, when assessing weight-loss and complications in 3 matched groups of nearly 3,000 patients each over 3 years [19].

As the low quality of evidence in data synthesis studies is a recurrent issue, there still exists a need for long-term data from a well-conducted, methodologically sound, pragmatic, multicentre, randomised trial of gastric banding and other bariatric techniques. Two on-going trials surgical randomised trials are the Swiss Multi-centre Bypass or Sleeve Study [20] and the UK By-Band randomised controlled trial [15]. In 2015 the latter has been converted to a three-arm trial of gastric banding, gastric bypass and sleeve gastrectomy (By-Band-Sleeve), and is the only such trial in the world comparing all three techniques.

Importantly these trials both include outcomes of weight and quality of life as primary outcome measures, and will also provide secondary outcome data on weight related co-morbidities and an economic evaluation of the study groups.

**Longer-term weight loss outcomes for gastric banding**

In the absence of well-conducted randomised trials with funded long-term data collection, long-term data of bariatric surgery are only available from observational studies and a few small, randomised trials. Most individual studies that quote >10 year data report that obstacles to follow-up impede the collection of accurate long-term data [21]. The longest follow up data is from the Swedish Obesity Study (SOS) [22], and has shown that bariatric surgery can reduce mortality, as well as cardiovascular and cancer risk over more than 10 years, in matched groups of over 2,000 patients. While two thirds of the patients in the SOS studies had a procedure that is no longer considered first line for weight loss (vertical banded gastroplasty), the study includes gastric bands with the longest follow up data being up to 20 years. The SOS shows maintenance of weight loss (18% of total body weight) in the all surgery groups at this time point, compared to 1% weight loss in controls, with sustainable reduction in diabetes rates, cardiovascular and cancer risks [23].

The only randomised trial with long term results of 10 years compared a small cohort (n = 51) of Italian patients randomised to gastric banding or gastric bypass [24]. The mean percent excess weight loss achieved at 10 years was 46 %  $\pm$  27 for gastric banding vs 69 %  $\pm$  29 for gastric bypass. With a mean BMI of patients in the two groups of 44, this equals a total body weight loss of 19 - 27%, comparable to that seen in the SOS study. A larger (n = 197) US randomised trial with 4 year follow up to 2007 measured similar excess weight loss outcomes with 45  $\pm$  28% for gastric band vs 68  $\pm$  19% for bypass in the medium-term but bypass was associated with more peri-operative and late complications and a higher early readmission rate [25]. A systematic review of randomised and prospective studies comparing gastric banding with gastric bypass, with more than 3-5 year follow up (29 studies

with 7971 patients), similarly showed that the mean sample size–weighted percentage of excess weight loss for gastric banding was 45.0% (n = 4109) vs 65.7% for gastric bypass (n = 3544) [26]. In all of these studies gastric band patients had a slower initial weight loss, which was typically maintained in the medium to long-term. In contrast, bypass patients lost weight rapidly in the first year and then started to regain some weight [23-26]. Additionally with gastric banding there is a proportion of patients, up to 15% who do not tolerate the band and fail to achieve any meaningful weight loss [27].

Gastric banding is the most predominant technique in Australia, and from there O'Brien et al have reported on 15-year follow up data from a large prospectively followed cohort of over 3,200 patients. The reported mean percent excess weight loss data is similar, 47% at 10 years (n = 714), and at 15 years (although in only 54 patients) [28]. This again equals 20.2% total body weight loss maintained at 10 - 15 years in the study population.

Other long-term studies are ongoing, such as the US Longitudinal Assessment of Bariatric Surgery [29], which is prospectively following nearly 2,500 patients after weight loss surgery, one quarter of whom had a gastric band. Three-year results published so far suggest average weight loss of 16% total body weight for those undergoing a gastric band, with most of the weight loss coming in the first year for all patients. However compared to gastric bypass patients (who started to regain some weight) the trajectory of weight change in the gastric banding group was maintained. In summary, all of the studies with at least 10 year follow up suggest that a clinically significant total body weight loss of 16 – 20 % is maintained in the long term following bariatric surgery, including gastric banding.

### **Diabetic outcomes with gastric banding**

Diabetes patients are an important group of those undergoing bariatric surgery, accounting for nearly one third of UK NHS bariatric patients [12, 13], and they have potentially the greatest benefits with a reduction in long-term diabetic complications. Early studies into bariatric surgery suggested that four fifths of all diabetic bariatric patients had a 'resolution'

of diabetes at 2 years following surgery [2930]. However a 2014 meta-analysis of longer term effects of metabolic surgery for type 2 diabetes, with stricter definitions for remission, reported a 65% remission rate and 89% remission or improvement rate in studies with more than 2 years follow up [3031]. The large UK national registry of bariatric surgery showed a 60 – 70% improvement in all diabetic patients at 3 years following surgery, with a decrease in medication use [12, 13].

A systematic review in 2012 of 35 studies involving gastric banding to treat diabetes showed remission or improvement in diabetes from 53% to 70% over 2 years, and although there was considerable heterogeneity in study quality the authors concluded that clinically relevant improvements in diabetes outcomes occurred in obese people with type 2 diabetes following LAGB surgery [3432]. The only randomised trial included in this review by Dixon et al, considered that participants randomized to surgical therapy were more likely to achieve remission of type 2 diabetes through greater weight loss [33]. The 2014 systematic review of studies comparing gastric banding with gastric bypass, included nine studies which measured co-morbidity improvement, and for type 2 diabetes (glycated haemoglobin <6.5% without medication), sample-size–weighted remission rates were 28.6% for gastric band (n = 96) and 66.7% for gastric bypass (n = 428) [26].

The interim analysis to assess the early control of type 2 diabetes (T2D), 1 year after gastric banding in the 5-year, prospective, observational Helping Evaluate Reduction in Obesity (HERO) study (N = 1,106) showed 72% achieved target control of diabetes compared with 42% at baseline [3234]. It is important to note that in these band studies, and many other studies of bariatric surgery, patients with shorter disease duration and not yet on insulin have the best likelihood of control of diabetes [12, 24, 30–31–3234]. Others have even performed studies of gastric banding in patients with lower BMIs and shown that remission of diabetes can be achieved in 50% of patients [3335]. Reflecting similar studies of all bariatric surgery in diabetics the 2014 UK NICE guidelines reduced the BMI threshold to 30 for ‘newly diagnosed diabetics’ to access bariatric surgery [10].

Bariatric surgery can also aid diabetes prevention. In an analysis of SOS patients without diabetes (n = 3,429) and a BMI of 34 – 38 at the start, after 15 years the incident rates of diabetes was 28.4 per 1,000 in the controls but only 6.8 per 1,000 in those who had surgery [3436].

### Complication rates

The major complications of gastric banding requiring re-operation or removal, are infection of the band, slippage and erosion, plus a group of patients who have intolerance of the band and fail to achieve any meaningful weight loss. For complications, the large systematic reviews of weight-loss surgery report reoperation rates as high as 40% [8, 9], but this includes some very early studies. Indeed, studies of bands placed pre-2000 reported rates of reoperation or removal as high as 60% [3537], and O'Brien's long term Australian study also reported similar revision rates of nearly 40% in the first 10 years after band placement [2728]. However, because of the evolution of band construction and operative technique, this had dropped to 6.4% over the last 5 years [2728].

The overall risks of all bariatric surgery itself are small, given the complex nature and co-morbidity burden of the patients involved, with the composite complication rate of gastric bypass (3.4%), being similar to that of laparoscopic cholecystectomy or hysterectomy [3638]. However, the reporting of adverse events in the outcomes of clinical trials is highly variable [3739]. A 2014 systematic review of the risks of bariatric surgery indicated a cumulative lifetime complication rate of 17% and a reoperation rate of 7%, but there were significant differences between the techniques [3827]. Thus, there were more early complications with gastric bypass than gastric banding, but the overall re-operation rate for banding was higher (although this mainly related to minor reoperations to the subcutaneous access port) [19, 24-2930]. Typically peri-operative complications occurred more frequently following gastric bypass than banding (8.0% vs 0.5%), while gastric bands had more long-term complications

1  
2  
3 requiring corrective procedures than gastric bypass (9% vs 2%). However, large volume  
4 gastric band units have shown that low re-operation rates, comparable overall to other  
5  
6 procedures long term, are achievable [2728]. Thus, it may be that at a population level, this  
7  
8 operation provides the greatest access to the benefits of weight loss surgery for the largest  
9  
10 number of patients.  
11

12  
13  
14 In view of this, it is surprising that the proportion of patients in the UK having gastric banding  
15  
16 has reduced over the last 5 years from 21% among NHS patients in 2009-10 to 14% in  
17  
18 2011-13 [12, 13]. Several factors may account for this: the rise in popularity of the gastric  
19  
20 sleeve operation (despite the paucity of long term data); the perceived difficulty with  
21  
22 arranging optimal long-term care for patients; and peer pressure among patients due to the  
23  
24 quicker weight loss attained with gastric bypass, even though weight loss at three years is  
25  
26 comparable between all 3 operations. In contrast, the proportion of self-funded patients  
27  
28 having banding was 42% in 2011-13, suggesting that it still maintains sufficient popularity  
29  
30 and attractiveness to patients, perhaps due to its affordability. Patients also commonly  
31  
32 perceive it as being less invasive than either gastric bypass or sleeve gastrectomy surgery, and  
33  
34 indeed it is technically less invasive than these stapling operations.  
35

36  
37 **Mortality rates**  
38

39  
40 The overall mortality rate for all primary bariatric surgery [12-14] is one tenth that of cardiac  
41  
42 surgery and comparable to elective arthroscopy, with the risk of gastric banding reportedly  
43  
44 being the lowest of the currently available surgical options (<0.01% vs 0.07% for gastric  
45  
46 bypass) [12, 19, 26, 3840]. Long term there is reportedly survival benefit compared to no  
47  
48 surgery for gastric banding patients [26, 2728].  
49

50  
51 **Revisional bariatric surgery and re-operations for gastric banding**  
52

53  
54 Bariatric surgery prevents and treats disease and disability, and improves quality of life, but  
55  
56 the need for reoperation (revisional surgery) reduces the clinical and cost effectiveness. In  
57  
58 addition to operations for complications (as above), revisional surgery may be considered for  
59  
60



later weight regain. However, the cumulative burden of revisional surgery on the patient and healthcare systems may be substantial, and revision surgery to the same or another bariatric operation (not including minor reoperations on the subcutaneous band access port) carries a 14 times increased mortality [12, 13, [3940](#)].

Weight regain after bariatric surgery is seen in a small but significant group of patients of all surgical techniques. Unfortunately it is not possible to predict before surgery which patients will fail to lose much weight or who will have later weight regain. A recent systematic review suggested multifactorial causes [[4041](#)], dietary, behavioural, physical, as well as surgical factors (in <20% of cases) which vary for the different techniques at different follow up times points. Clearly, for gastric banding, intensive input from all members of the MDT, and frequent (monthly) follow-up intervals after surgery are particularly important in producing and maintaining weight-loss [[4142](#) - [4344](#)]. As dietary, psychological and physical inactivity were the main factors in up to 80% of weight regain, addressing these requires a clinical service with a systematic approach to assess and identify those patients who may have a surgical factor for weight regain which might benefit from revisional surgery. Importantly, current data suggest that every patient who has had a gastric band removed, for any reason, will have weight regain back to or near their baseline level [[4445](#)].

### **Cost effectiveness of gastric banding**

The long term benefits of bariatric surgery for weight loss, co-morbidity resolution, lower cancer rates and improved mortality could be expected to lead to a health care cost saving but results are not that clear cut. While a systematic review in 2011 [[4546](#)] suggested that weight-loss surgery was cost effective compared to non-surgical treatment, a 2014 study on insurance provider claim costs after bariatric surgery showed no reduction in actual costs of claims [[4647](#)]. However, the measure used for comparison is important, and many studies in different countries have shown that the incremental cost effectiveness ratio per quality- or disability- adjusted life year varied from about £1,300 for gastric banding in Australia [[4847](#)],



to £4,000 - £4,500 for all laparoscopic weight loss surgery in the USA [4948]. Interestingly, while all these values are far below the cost-effectiveness threshold for the respective countries (and also for the UK), gastric banding may be the most cost saving treatment at a population level.

An analysis of data using a decision analytic model from the Scandinavian Registry, covering cardiovascular disease, diabetes and surgical complications, showed that bariatric surgery led to a cost saving when compared to non-surgical management of weight loss [5049]. When the risk reduction of vascular events, reduction in diabetes and gain in quality adjusted life years were extrapolated for the whole cohort, operated on in one year, the lifetime gains were over 32,000 quality-adjusted person years, and a net saving of £43 million in health care costs.

The very low volume of bariatric surgery in the UK represents much less than 1 in 100 of the 5%, or 2.1 million people [5051], who meet UK NICE guidelines for surgery. In other developed countries ~~as well~~ there are large differences in the prevalence of co-morbidities and socio-demographic status between surgery-eligible patients and bariatric surgery recipients [5254]. Clearly no healthcare system could provide capacity to treat such a potentially great demand, but the overwhelming evidence is that surgery is clinically beneficial, and cost effective. Thus, the RCT evidence in the NICE Guidance showed clear improvement in every aspect of diabetes control and on this basis increasing availability of bariatric surgery should be a clear aspiration for healthcare services. Alternatively, the availability of government funded weight-loss surgery could be limited to the procedure with the lowest costs and shortest hospitalization requirements, to maximise the opportunity for delivery of a service. This strategy would favour gastric banding, although this does not take into account the longer term follow up costs, and the provision of services required, for gastric band adjustments in the community or specialised clinics.

**CONCLUSIONS**

Obesity, with nearly 30 million sufferers in the UK currently, and its treatment, is a huge problem for the NHS. Current evidence shows that only bariatric surgery gives clinically meaningful weight loss, although there is a paucity of high quality long-term evidence. Despite its effectiveness less than 5,000 bariatric procedures were performed in the NHS in 2013 [12, 13], thus the NHS provides the life-saving and quality of life improving benefit of surgery to less than 1 in 4,000 of the severely obese population in the UK. There would need to be a huge turn-around in obesity service provision for any significant population health gains to be achieved. The nature of which operation would best deliver these gains is still unknown, more high quality studies are needed, but with the lowest reported incremental cost effectiveness ratio per quality adjusted life year, gastric banding may provide the NHS with the most punch for its pound. However, unlike Australia, the UK NHS is not currently well set up to look after large number of gastric band patients and adjust the bands regularly in hospital (outside of clinical trials) or in the community. This issue of gastric band follow-up is probably the reason why many patients and surgeons do not choose gastric banding over other operations in the NHS. In addition, many bands are removed by surgeons in non-specialist centres if the patient develops a problem such as slippage, rather than having it rectified. It would also require a shift in the moral perspective of delivering treatments effective at improving quality of life to those with a significant need [5253]. Still a major unknown remains the costs associated with the management of the long-term complications of bariatric surgery, including that of modern gastric banding, which appear comparable for all techniques.

References

1. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384(9945):766-81.

2. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes research and clinical practice*. 2014;103(2):137-49.

3. De Pergola G, Silvestris F. Obesity as a major risk factor for cancer. *Journal of obesity*. 2013;2013:291546.

4. Herrigel DJ, Moss RA. Diabetes mellitus as a novel risk factor for gastrointestinal malignancies. *Postgraduate medicine*. 2014;126(6):106-18.

5. Korhonen PE, Seppälä T, Järvenpää S, Kautiainen H. Body mass index and health-related quality of life in apparently healthy individuals. *Qual Life Res*. 2014 Feb;23(1):67-74.

6. Adams TD, Davidson LE, Litwin SE, Kolotkin RL, LaMonte MJ, Pendleton RC et al. Health benefits of gastric bypass surgery after 6 years. *Jama*. 2012;308(11):1122-31.

7. Mehta T, Fontaine KR, Keith SW, Bangalore SS, de los Campos G, Bartolucci A et al. Obesity and mortality: are the risks declining? Evidence from multiple prospective studies in the United States. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2014;15(8):619-29.

8. Colquitt JL, Pickett K, Loveman E, Frampton GK. Surgery for weight loss in adults. *The Cochrane database of systematic reviews*. 2014;8:Cd003641.

9. Gloy VL, Briel M, Bhatt DL, Kashyap SR, Schauer PR, Mingrone G et al. Bariatric surgery versus non-surgical treatment for obesity: a systematic review and meta-analysis of randomised controlled trials. *BMJ (Clinical research ed)*. 2013;347:f5934.

10. National Clinical Guideline CG189. National Institute for Health and Clinical Excellence: Guidance. Obesity: Identification, Assessment and Management of Overweight and Obesity in Children, Young People and Adults: Partial Update of CG43. London: National Institute for Health and Care Excellence (UK) Copyright (c) National Clinical Guideline Centre, 2014.

11. Buchwald H, Oien DM. Metabolic/bariatric surgery worldwide 2011. *Obesity surgery*. 2013;23(4):427-36.

12. Welbourn R, Small P, Finlay I, Sareela A, Somers S, Mahawar K et al. The Second National Bariatric Surgery Registry Report to March 2013. Dendrite Clinical Systems; Henley UK. ISBN 978-0-9568154-8-4 (2014).

13. Welbourn R, Fiennes A, Kinsman R, Walton P. The UK National Bariatric Surgery Registry. First Report to 2010. Dendrite Clinical Systems, Henley UK. ISBN 1-903968-27-5 (2011).

14. BOMSS. Consultant Outcomes 2013. 2013. <http://www.bomss.org.uk/surgeon-data/>.
15. Rogers CA, Welbourn R, Byrne J, Donovan JL, Reeves BC, Wordsworth S et al. The By-Band study: gastric bypass or adjustable gastric band surgery to treat morbid obesity: study protocol for a multi-centre randomised controlled trial with an internal pilot phase. *Trials*. 2014;15:53.
16. Burton PR, Yap K, Brown WA, Laurie C, O'Donnell M, Hebbard G, Kalff V, O'Brien PE. Changes in satiety, supra- and infraband transit, and gastric emptying following laparoscopic adjustable gastric banding: a prospective follow-up study. *Obes Surg*. 2011 Feb;21(2):217-23.
17. Moore M, Wainwright P, Hopkins JC. Practice pointer: Primary care management of patients after weight loss surgery. *BMJ* in press
18. Brown W, Korin A, Burton P, O'Brien PE. Laparoscopic adjustable gastric banding. *Aust Fam Physician*. 2009 Dec;38(12):972-6.
19. Carlin AM, Zeni TM, English WJ, Hawasli AA, Genaw JA, Krause KR et al. The comparative effectiveness of sleeve gastrectomy, gastric bypass, and adjustable gastric banding procedures for the treatment of morbid obesity. *Annals of surgery*. 2013;257(5):791-7.
20. Peterli R, Borbely Y, Kern B, Gass M, Peters T, Thurnheer M et al. Early results of the Swiss Multicentre Bypass or Sleeve Study (SM-BOSS): a prospective randomized trial comparing laparoscopic sleeve gastrectomy and Roux-en-Y gastric bypass. *Annals of surgery*. 2013;258(5):690-4
21. Higa K, Ho T, Tercero F, Yunus T, Boone KB. Laparoscopic Roux-en-Y gastric bypass: 10-year follow-up. *Surgery for obesity and related diseases*. 2011;7(4):516-25.
22. Sjostrom L, Narbro K, Sjostrom CD, Karason K, Larsson B, Wedel H et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *The New England journal of medicine*. 2007;357(8):741-52.
23. Sjostrom L. Review of the key results from the Swedish Obese Subjects (SOS) trial - a prospective controlled intervention study of bariatric surgery. *Journal of internal medicine*. 2013;273(3):219-34.
24. Angrisani L, Cutolo PP, Formisano G, Nosso G, Vitolo G. Laparoscopic adjustable gastric banding versus Roux-en-Y gastric bypass: 10-year results of a prospective, randomized trial. *Surgery for obesity and related diseases*. 2013;9(3):405-13.
25. Nguyen NT, Slone JA, Nguyen XM, Hartman JS, Hoyt DB. A prospective randomized trial of laparoscopic gastric bypass versus laparoscopic adjustable gastric banding for the treatment of morbid obesity: outcomes, quality of life, and costs. *Ann Surg*. 2009 Oct;250(4):631-41.
26. Puzziferri N, Roshek TB 3rd, Mayo HG, Gallagher R, Belle SH, Livingston EH. Long-term follow-up after bariatric surgery: a systematic review. *JAMA*. 2014 Sep 3;312(9):934-42.
27. [Chang S, Stoll CT, Song J, Varela J, Eagon CJ, Colditz GA. The effectiveness and risks of bariatric surgery: An updated systematic review and meta-analysis, 2003-2012. \*JAMA Surgery\*. 2014;149\(3\):275-87.](#)

28. O'Brien PE, MacDonald L, Anderson M, Brennan L, Brown WA. Long-term outcomes after bariatric surgery: fifteen-year follow-up of adjustable gastric banding and a systematic review of the bariatric surgical literature. *Annals of surgery*. 2013;257(1):87-94.

289. Courcoulas AP, Christian NJ, Belle SH, Berk PD, Flum DR, Garcia L et al. Weight change and health outcomes at 3 years after bariatric surgery among individuals with severe obesity. *Jama*. 2013;310(22):2416-25.

3029. Buchwald H, Estok R, Fahrbach K, Banel D, Jensen MD, Pories WJ et al. Weight and type 2 diabetes after bariatric surgery: systematic review and meta-analysis. *The American journal of medicine*. 2009;122(3):248-56.e5.

3031. Yu J, Zhou X, Li L, Li S, Tan J, Li Y et al. The long-term effects of bariatric surgery for type 2 diabetes: systematic review and meta-analysis of randomized and non-randomized evidence. *Obesity surgery*. 2015;25(1):143-58.

3132. Dixon JB, Murphy DK, Segel JE, Finkelstein EA. Impact of laparoscopic adjustable gastric banding on type 2 diabetes. *Obes Rev*. 2012 Jan;13(1):57-67.

3233. Dixon JB, O'Brien PE, Playfair J, Chapman L, Schachter LM, Skinner S, Proietto J, Bailey M, Anderson M. Adjustable gastric banding and conventional therapy for type 2 diabetes: a randomized controlled trial. *JAMA*. 2008 Jan 23;299(3):316-23. doi: 10.1001/jama.299.3.316.

34. Edelman S, Ng-Mak DS, Fusco M, Ashton D, Okerson T, Liu Q, Jin J, Dixon JB. Control of type 2 diabetes after 1 year of laparoscopic adjustable gastric banding in the helping evaluate reduction in obesity (HERO) study. *Diabetes Obes Metab*. 2014 Oct;16(10):1009-15

3335. Wentworth JM, Playfair J, Laurie C, Ritchie ME, Brown WA, Burton P, Shaw JE, O'Brien PE. Multidisciplinary diabetes care with and without bariatric surgery in overweight people: a randomised controlled trial. *Lancet Diabetes Endocrinol*. 2014 Jul;2(7):545-52.

3436. Carlsson LM, Peltonen M, Ahlin S, Anveden A, Bouchard C, Carlsson B et al. Bariatric surgery and prevention of type 2 diabetes in Swedish obese subjects. *The New England journal of medicine*. 2012;367(8):695-704.

3537. Himpens J, Cadière GB, Bazi M, Vouche M, Cadière B, Dapri G. Long-term outcomes of laparoscopic adjustable gastric banding. *Arch Surg*. 2011 Jul;146(7):802-7.

3638. Aminian A, Brethauer SA, Kirwan JP, Kashyap SR, Burguera B, Schauer PR. How safe is metabolic/diabetes surgery? *Diabetes, obesity & metabolism*. 2015;17(2):198-201.

3739. Hopkins JC, Howes N, Chalmers K, Savovic J, Whale K, Coulman KD et al. Outcome reporting in bariatric surgery: an in-depth analysis to inform the development of a core outcome set, the BARIACT Study. *Obesity reviews*. 2015;16(1):88-106.

38. Chang S, Stoll CT, Song J, Varela J, Eagon CJ, Colditz GA. The effectiveness and risks of bariatric surgery: An updated systematic review and meta-analysis, 2003-2012. *JAMA Surgery*. 2014;149(3):275-87.

- [3940](#). Sudan R, Nguyen NT, Hutter MM, Brethauer SA, Ponce J, Morton JM. Morbidity, mortality, and weight loss outcomes after reoperative bariatric surgery in the USA. *Journal of gastrointestinal surgery*. 2015;19(1):171-8; discussion 8-9.
- [4041](#). Karmali S, Brar B, Shi X, Sharma AM, de Gara C, Birch DW. Weight recidivism post-bariatric surgery: a systematic review. *Obesity surgery*. 2013;23(11):1922-33.
- [4142](#). te Riele WW, Boerma D, Wiezer MJ, Borel Rinkes IH, van Ramshorst B. Long-term results of laparoscopic adjustable gastric banding in patients lost to follow-up. *Br J Surg*. 2010 Oct;97(10):1535-40.
- [4243](#). Hochberg LS, Murphy KD, O'Brien PE, Brennan L. Laparoscopic Adjustable Gastric Banding (LAGB) Aftercare Attendance and Attrition. *Obes Surg*. 2015 Sep;25(9):1693-702.
- [4344](#). Sysko R, Hildebrandt TB, Kaplan S, Brewer SK, Zitsman JL, Devlin MJ. Predictors and correlates of follow-up visit adherence among adolescents receiving laparoscopic adjustable gastric banding. *Surg Obes Relat Dis*. 2014 Sep-Oct;10(5):914-20.
- [4445](#). Aarts EO, Dogan K, Koehestanie P, Janssen IM, Berends FJ. What happens after gastric band removal without additional bariatric surgery? *Surg Obes Relat Dis*. 2014 Nov-Dec;10(6):1092-6.
- [4546](#). Padwal R, Klarenbach S, Wiebe N, Hazel M, Birch D, Karmali S et al. Bariatric surgery: a systematic review of the clinical and economic evidence. *Journal of general internal medicine*. 2011;26(10):1183-94.
- [4647](#). Weiner JP, Goodwin SM, Chang HY, Bolen SD, Richards TM, Johns RA et al. Impact of bariatric surgery on health care costs of obese persons: a 6-year follow-up of surgical and comparison cohorts using health plan data. *JAMA Surg*. 2013;148(6):555-62.
- [4748](#). Lee YY, Veerman JL, Barendregt JJ. The cost-effectiveness of laparoscopic adjustable gastric banding in the morbidly obese adult population of Australia. *PloS one*. 2013;8(5):e64965.
- [4849](#). Wang BC, Wong ES, Alfonso-Cristancho R, He H, Flum DR, Arterburn DE et al. Cost-effectiveness of bariatric surgical procedures for the treatment of severe obesity. *The European journal of health economics:HEPAC* 2014;15(3):253-63.
- [4950](#). Borisenko O, Adam D, Funch-Jensen P, Ahmed AR, Zhang R, Colpan Z et al. Bariatric Surgery can Lead to Net Cost Savings to Health Care Systems: Results from a Comprehensive European Decision Analytic Model. *Obesity surgery*. 2015 Sep;25(9):1559-68.
- [5051](#). Ahmad A, Laverty AA, Aasheim E, Majeed A, Millett C, Saxena S. Eligibility for bariatric surgery among adults in England: analysis of a national cross-sectional survey. *JRSM Open*. 2014 Jan 7;5(1):2042533313512479. doi: 10.1177/2042533313512479. eCollection 2014.
- [5152](#). Padwal RS, Chang HJ, Klarenbach S, Sharma AM, Majumdar SR. Characteristics of the population eligible for and receiving publicly funded bariatric surgery in Canada. *International journal for equity in health*. 2012;11:54.
- [5253](#). Persson K. Why Bariatric surgery should be given high priority: an argument from law and morality. *Health care analysis:HCA: journal of health philosophy and policy*. 2014;22(4):305-24.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

# The use of adjustable gastric bands for management of severe and complex obesity

James C A Hopkins DPhil FRCS<sup>1</sup>,

Jane M Blazeby MD FRCS<sup>2</sup>

Chris A Rogers PhD<sup>3</sup>

Richard Welbourn MD FRCS<sup>1</sup>

<sup>1</sup> Department of Upper Gastrointestinal Surgery and Bariatric Surgery, Musgrove Park Hospital, Taunton TA1 5DA, UK

<sup>2</sup> Centre for Surgical Research, School of Social and Community Medicine, University of Bristol, Bristol

<sup>3</sup> Clinical Trials and Evaluation Unit, School of Clinical Sciences, University of Bristol, Bristol, UK

**Number of Figures = 1**

**Number of Colour Figures = 1**

**Number of Tables = 0**

Figure 1 has been previously published and permission has been given from Elsevier.

There is no supplementary material to go online.

**Running title:** Adjustable gastric banding for severe and complex obesity

**Keywords:** Obesity surgery, bariatric surgery, diabetes treatment

**Author for correspondence:** Mr Richard Welbourn, Consultant Surgeon, Dept Upper GI and Bariatric Surgery, Musgrove Park Hospital, Taunton TA1 5DA, UK



Richard.Welbourn@tst.nhs.uk, Tel +44 (0)1823 342100, Fax +44 (0)1823 343560

**Other authors' details:**

JCA Hopkins, Fellow in Upper GI and Bariatric Surgery, frog@doctors.org.uk

JM Blazeby, Professor of Surgery, J.M.blazeby@bristol.ac.uk

CA Rogers, Reader in Medical Statistics, Chris.Rogers@bristol.ac.uk

RW acts as guarantor and confirms that the manuscript has been submitted solely to this journal and is not published, in press, or submitted elsewhere.

**Conflict of Interest declaration:** James Hopkins, Jane Blazeby and Chris Rogers declare no conflict of interest. Richard Welbourn received sponsorship for attending conference/educational workshops from Allergan and Ethicon Endo-Surgery and is a consultant for Novo Nordisk. Musgrove Park Hospital, Taunton UK, also received grant funding for a bariatric surgery training fellowship from Ethicon Endo-Surgery.

Key words: obesity surgery, bariatric surgery, metabolic surgery, gastric banding

## ABSTRACT

### Background

Obesity levels in the UK have reached a sustained high and about 5% of the population would be candidates for bariatric surgery based upon current UK NICE guidelines, which has important implications for Clinical Commissioning Groups.

### Sources of data

Summary data from Cochrane systematic reviews, randomised controlled trials (RCTs) and cohort studies.

### Areas of agreement

Currently the only treatment that offers significant and durable weight-loss for those with severe and complex obesity is surgery. Three operations account for 95% of all bariatric surgery in the UK, but the NHS offers surgery to only a small fraction of those who could benefit. Laparoscopic adjustable gastric banding (gastric banding) has potentially the lowest risk and up-front costs of the three procedures.

### Areas of controversy

Reliable level 1 evidence of the relative effectiveness of the operations is lacking.

### Growing points

As a point intervention, weight-loss surgery together with the chronic disease management strategy for obesity can prevent significant future disease and mortality, and the NHS should embrace both.

### Areas timely for developing research

Better RCT evidence is needed including clinical effectiveness and economic analysis to answer the important question 'which is the best of the three operations most frequently

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

performed?’ This review considers the current evidence for gastric banding for the treatment of severe and complex obesity.

For Peer Review

## INTRODUCTION

Obesity is a major worldwide public health problem with significant implications for primary care, particularly where obesity prevalence in developed countries is as high as 32.5% in women in high income North American countries, 29.8% in Australasia and 21.0% in Western Europe [1]. Although obesity levels may have reached a plateau, the burden of the important obesity related co-morbidities, such as type 2 diabetes, which was 382 million people worldwide in 2013, is predicted to rise to over 590 million in 2035 [2]. Both diabetes and obesity are independent risk factors for many important cancers [3, 4]. Obesity significantly increases many functional disorders and worsens overall health related quality of life [5], which can be improved by weight loss and metabolic surgery [6]. Overall, there is substantial evidence that obesity is a chronic disease that shortens life expectancy, even with the current best medical management of its co-morbidities [7].

Bariatric surgery is the only reproducibly effective treatment for severe and complex obesity [8, 9]. UK NICE guidelines provide clear guidance for the delivery of health-care and recommend that: bariatric surgery is the first line of treatment for those with a BMI of  $\geq 50$  in whom lifestyle interventions have failed; and those with a BMI  $\geq 40$ , or  $\geq 35$  with severe obesity-related co-morbidities, should be offered bariatric surgery [10]. Weight-loss and metabolic surgery procedures are among the most commonly performed elective general surgical operations in the world, with over 350,000 performed in 2011, [11] but in 2013 in the UK, with government funded health care, only around 8,000 operations were performed [12, 13], of which around 4,300 operations are recorded as being performed in the public sector [14].

Bariatric surgery is a young surgical speciality, with increasing numbers of cases being performed over a relatively short time frame [11], and there is a lack of high quality long term outcome data for the procedures. The most up-to-date, 2014, Cochrane collaboration systematic review of weight loss surgery in adults could only provide synthesizable evidence


for 3 years of follow up, and commented that ‘the long-term effects of surgery remain unclear’ [8]. However this level 1 evidence for short and medium term results of surgery can be compared directly to results for non-surgical interventions, and combined with some long-term cohort studies of bariatric surgery, albeit in smaller numbers, to show a pattern of long-term benefits from surgery. Given the established long-term benefit it is important to have large scale, pragmatic randomised controlled surgical trials to evaluate different techniques, to include economic evaluations, to give us the answer as to what is the best operation [15].

**Gastric banding and how it facilitates weight loss**

In the UK >99% of gastric band procedures are performed laparoscopically and patients typically spend ≤1 days in hospital with very few early complications [12, 13]. The operation induces and sustains weight loss by activating a satiety mechanism. Adding or removing fluid from the band achieves optimal filling. Food is not retained above the optimally adjusted band, i.e. not physically restricted, but briefly delayed food bolus transit through the band and the continuous pressure of the optimally filled band on the stomach wall produces both early satiety and a lack of appetite [16]. The effects of the band on oesophageal and proximal gastric function appears to activate a satiety signal, transmitted to CNS satiety centres via the vagus nerve, without physically restricting the meal size [16]. Figure 1 shows the position of a gastric band with a small “virtual” pouch of stomach below the gastro-oesophageal junction, above the final band position, held within gastro-gastro tunnelling sutures. Ideally patients are followed up every 1 – 3 months for the first two years by the bariatric multidisciplinary team (MDT), and then yearly in conjunction with GPs as part of a shared care model of chronic disease management (NICE CG189) [10].

Patients require life-long follow up for optimal weight control and co-morbidity resolution for gastric banding to work, especially as nutritional issues or complications can occur at any point after surgery [17, 18]. Symptoms of reflux, regurgitation or vomiting and especially nocturnal aspiration or dysphagia should alert the primary care GP to possible

complications. Abdominal pain along with any of these symptoms may suggest an acute complication, but other causes of abdominal pain should always be considered. Lifelong care involves essential counselling about food choices and eating patterns, as well as adjustment of the band [18]. Patients should therefore not be offered revisional surgery unless a process for continuing care is in place with the bariatric multidisciplinary team and within primary care.



**Figure 1** The adjustable gastric band. (Images reproduced from SM Griffin, SA Raimes, J Shenfine. *Oesophagogastric Surgery*. 5th ed. Saunders Elsevier, 2013)

### Weight loss effects of gastric bands & bariatric surgery

The 2014 Cochrane review into weight loss surgery for adults included 22 randomised trials with 1,798 participants. The authors compared surgery with non-surgical interventions or different surgical procedures and most studies followed participants for only 12 to 36 months [8]. All of the trials which compared surgery with non-surgical interventions for weight loss found that surgery was significantly superior for weight loss at 1-2 years. This mirrored a 2013 systematic review and meta-analysis that directly compared surgery and medical therapy, using 11 randomised studies with 796 included patients, and a random effect model, clearly favouring surgery [9]. With study follow up of up to 2 years individuals

undergoing bariatric surgery lost more body weight (mean difference -26 kg (95% confidence interval -31 to -21),  $P<0.001$ ) than those with medical therapy alone.

In the Cochrane review no clinically significant differences between the three main techniques of gastric band, all types of gastric bypass or sleeve gastrectomy existed for the majority of time points within 'multi-trial' evidence. Although only three trials directly compared gastric banding with laparoscopic Roux-en-Y gastric bypass (gastric bypass), the latter produced greater weight loss, with a mean difference in body mass index (BMI) reduction up to five years of -5.2 kg/m<sup>2</sup> (95% confidence interval -6.4 to -4.0;  $P<0.00001$ ).

However, this was only from a total of 265 patients when combined, and the quality of the evidence was described as *moderate* at best. In the seven included trials of gastric bypass and sleeve gastrectomy there was no consistent picture as to which procedure was better or worse, but with a *low* quality of evidence [8]. For the 2013 systematic review the cumulative body weight loss was similar but not statistically significantly different between gastric banding and gastric bypass or other techniques combined (mean difference between 6 and 7 kg,  $P>0.20$ ) [9]. Effects on co-morbidities, complications and additional surgical procedures were neutral for the three main procedures in these reviews [8, 9]. Similarly a large prospective database, the Michigan Bariatric Surgery Collaborative, concluded that there was comparable effectiveness of gastric banding, gastric bypass and sleeve gastrectomy for the treatment of severe obesity, when assessing weight-loss and complications in 3 matched groups of nearly 3,000 patients each over 3 years [19].

As the low quality of evidence in data synthesis studies is a recurrent issue, there still exists a need for long-term data from a well-conducted, methodologically sound, pragmatic, multicentre, randomised trial of gastric banding and other bariatric techniques. Two on-going trials surgical randomised trials are the Swiss Multi-centre Bypass or Sleeve Study [20] and the UK By-Band randomised controlled trial [15]. In 2015 the latter has been converted to a three-arm trial of gastric banding, gastric bypass and sleeve gastrectomy (By-Band-Sleeve), and is the only such trial in the world comparing all three techniques.

Importantly these trials both include outcomes of weight and quality of life as primary outcome measures, and will also provide secondary outcome data on weight related co-morbidities and an economic evaluation of the study groups.

### **Longer-term weight loss outcomes for gastric banding**

In the absence of well-conducted randomised trials with funded long-term data collection, long-term data of bariatric surgery are only available from observational studies and a few small, randomised trials. Most individual studies that quote >10 year data report that obstacles to follow-up impede the collection of accurate long-term data [21]. The longest follow up data is from the Swedish Obesity Study (SOS) [22], and has shown that bariatric surgery can reduce mortality, as well as cardiovascular and cancer risk over more than 10 years, in matched groups of over 2,000 patients. While two thirds of the patients in the SOS studies had a procedure that is no longer considered first line for weight loss (vertical banded gastroplasty), the study includes gastric bands with the longest follow up data being up to 20 years. The SOS shows maintenance of weight loss (18% of total body weight) in the all surgery groups at this time point, compared to 1% weight loss in controls, with sustainable reduction in diabetes rates, cardiovascular and cancer risks [23].

The only randomised trial with long term results of 10 years compared a small cohort (n = 51) of Italian patients randomised to gastric banding or gastric bypass [24]. The mean percent excess weight loss achieved at 10 years was 46 %  $\pm$  27 for gastric banding vs 69 %  $\pm$  29 for gastric bypass. With a mean BMI of patients in the two groups of 44, this equals a total body weight loss of 19 - 27%, comparable to that seen in the SOS study. A larger (n = 197) US randomised trial with 4 year follow up to 2007 measured similar excess weight loss outcomes with 45  $\pm$  28% for gastric band vs 68  $\pm$  19% for bypass in the medium-term but bypass was associated with more peri-operative and late complications and a higher early readmission rate [25]. A systematic review of randomised and prospective studies comparing gastric banding with gastric bypass, with more than 3-5 year follow up (29 studies



with 7971 patients), similarly showed that the mean sample size–weighted percentage of excess weight loss for gastric banding was 45.0% (n = 4109) vs 65.7% for gastric bypass (n = 3544) [26]. In all of these studies gastric band patients had a slower initial weight loss, which was typically maintained in the medium to long-term. In contrast, bypass patients lost weight rapidly in the first year and then started to regain some weight [23-26]. Additionally with gastric banding there is a proportion of patients, up to 15% who do not tolerate the band and fail to achieve any meaningful weight loss [27].

Gastric banding is the most predominant technique in Australia, and from there O'Brien et al have reported on 15-year follow up data from a large prospectively followed cohort of over 3,200 patients. The reported mean percent excess weight loss data is similar, 47% at 10 years (n = 714), and at 15 years (although in only 54 patients) [28]. This again equals 20.2% total body weight loss maintained at 10 - 15 years in the study population.

Other long-term studies are ongoing, such as the US Longitudinal Assessment of Bariatric Surgery [29], which is prospectively following nearly 2,500 patients after weight loss surgery, one quarter of whom had a gastric band. Three-year results published so far suggest average weight loss of 16% total body weight for those undergoing a gastric band, with most of the weight loss coming in the first year for all patients. However compared to gastric bypass patients (who started to regain some weight) the trajectory of weight change in the gastric banding group was maintained. In summary, all of the studies with at least 10 year follow up suggest that a clinically significant total body weight loss of 16 – 20 % is maintained in the long term following bariatric surgery, including gastric banding.

**Diabetic outcomes with gastric banding**

Diabetes patients are an important group of those undergoing bariatric surgery, accounting for nearly one third of UK NHS bariatric patients [12, 13], and they have potentially the greatest benefits with a reduction in long-term diabetic complications. Early studies into bariatric surgery suggested that four fifths of all diabetic bariatric patients had a 'resolution'

of diabetes at 2 years following surgery [30]. However a 2014 meta-analysis of longer term effects of metabolic surgery for type 2 diabetes, with stricter definitions for remission, reported a 65% remission rate and 89% remission or improvement rate in studies with more than 2 years follow up [31]. The large UK national registry of bariatric surgery showed a 60 – 70% improvement in all diabetic patients at 3 years following surgery, with a decrease in medication use [12, 13].

A systematic review in 2012 of 35 studies involving gastric banding to treat diabetes showed remission or improvement in diabetes from 53% to 70% over 2 years, and although there was considerable heterogeneity in study quality the authors concluded that clinically relevant improvements in diabetes outcomes occurred in obese people with type 2 diabetes following LAGB surgery [32]. The only randomised trial included in this review by Dixon et al, considered that participants randomized to surgical therapy were more likely to achieve remission of type 2 diabetes through greater weight loss [33]. The 2014 systematic review of studies comparing gastric banding with gastric bypass, included nine studies which measured co-morbidity improvement, and for type 2 diabetes (glycated haemoglobin <6.5% without medication), sample-size-weighted remission rates were 28.6% for gastric band (n = 96) and 66.7% for gastric bypass (n = 428) [26].

The interim analysis to assess the early control of type 2 diabetes (T2D), 1 year after gastric banding in the 5-year, prospective, observational Helping Evaluate Reduction in Obesity (HERO) study (N = 1,106) showed 72% achieved target control of diabetes compared with 42% at baseline [34]. It is important to note that in these band studies, and many other studies of bariatric surgery, patients with shorter disease duration and not yet on insulin have the best likelihood of control of diabetes [12, 24, 31-34]. Others have even performed studies of gastric banding in patients with lower BMIs and shown that remission of diabetes can be achieved in 50% of patients [35]. Reflecting similar studies of all bariatric surgery in diabetics the 2014 UK NICE guidelines reduced the BMI threshold to 30 for 'newly diagnosed diabetics' to access bariatric surgery [10].

Bariatric surgery can also aid diabetes prevention. In an analysis of SOS patients without diabetes (n = 3,429) and a BMI of 34 – 38 at the start, after 15 years the incident rates of diabetes was 28.4 per 1,000 in the controls but only 6.8 per 1,000 in those who had surgery [36].

**Complication rates**

The major complications of gastric banding requiring re-operation or removal, are infection of the band, slippage and erosion, plus a group of patients who have intolerance of the band and fail to achieve any meaningful weight loss. For complications, the large systematic reviews of weight-loss surgery report reoperation rates as high as 40% [8, 9], but this includes some very early studies. Indeed, studies of bands placed pre-2000 reported rates of reoperation or removal as high as 60% [37], and O'Brien's long term Australian study also reported similar revision rates of nearly 40% in the first 10 years after band placement [28]. However, because of the evolution of band construction and operative technique, this had dropped to 6.4% over the last 5 years [28].

The overall risks of all bariatric surgery itself are small, given the complex nature and co-morbidity burden of the patients involved, with the composite complication rate of gastric bypass (3.4%), being similar to that of laparoscopic cholecystectomy or hysterectomy [38]. However, the reporting of adverse events in the outcomes of clinical trials is highly variable [39]. A 2014 systematic review of the risks of bariatric surgery indicated a cumulative lifetime complication rate of 17% and a reoperation rate of 7%, but there were significant differences between the techniques [27]. Thus, there were more early complications with gastric bypass than gastric banding, but the overall re-operation rate for banding was higher (although this mainly related to minor reoperations to the subcutaneous access port) [19, 24-30]. Typically peri-operative complications occurred more frequently following gastric bypass than banding (8.0% vs 0.5%), while gastric bands had more long-term complications requiring corrective

procedures than gastric bypass (9% vs 2%). However, large volume gastric band units have shown that low re-operation rates, comparable overall to other procedures long term, are achievable [28]. Thus, it may be that at a population level, this operation provides the greatest access to the benefits of weight loss surgery for the largest number of patients.

In view of this, it is surprising that the proportion of patients in the UK having gastric banding has reduced over the last 5 years from 21% among NHS patients in 2009-10 to 14% in 2011-13 [12, 13]. Several factors may account for this: the rise in popularity of the gastric sleeve operation (despite the paucity of long term data); the perceived difficulty with arranging optimal long-term care for patients; and peer pressure among patients due to the quicker weight loss attained with gastric bypass, even though weight loss at three years is comparable between all 3 operations. In contrast, the proportion of self-funded patients having banding was 42% in 2011-13, suggesting that it still maintains sufficient popularity and attractiveness to patients, perhaps due to its affordability. Patients also commonly perceive it as being less invasive than either gastric bypass or sleeve gastrectomy surgery, and indeed it is technically less invasive than these stapling operations.

### **Mortality rates**

The overall mortality rate for all primary bariatric surgery [12-14] is one tenth that of cardiac surgery and comparable to elective arthroscopy, with the risk of gastric banding reportedly being the lowest of the currently available surgical options (<0.01% vs 0.07% for gastric bypass) [12, 19, 26, 40]. Long term there is reportedly survival benefit compared to no surgery for gastric banding patients [26, 28].

### **Revisional bariatric surgery and re-operations for gastric banding**

Bariatric surgery prevents and treats disease and disability, and improves quality of life, but the need for reoperation (revisional surgery) reduces the clinical and cost effectiveness. In addition to operations for complications (as above), revisional surgery may be considered for later weight regain. However, the cumulative burden of revisional surgery on the patient and

healthcare systems may be substantial, and revision surgery to the same or another bariatric operation (not including minor reoperations on the subcutaneous band access port) carries a 14 times increased mortality [12, 13, 40].

Weight regain after bariatric surgery is seen in a small but significant group of patients of all surgical techniques. Unfortunately it is not possible to predict before surgery which patients will fail to lose much weight or who will have later weight regain. A recent systematic review suggested multifactorial causes [41], dietary, behavioural, physical, as well as surgical factors (in <20% of cases) which vary for the different techniques at different follow up times points. Clearly, for gastric banding, intensive input from all members of the MDT, and frequent (monthly) follow-up intervals after surgery are particularly important in producing and maintaining weight-loss [42 - 44]. As dietary, psychological and physical inactivity were the main factors in up to 80% of weight regain, addressing these requires a clinical service with a systematic approach to assess and identify those patients who may have a surgical factor for weight regain which might benefit from revisional surgery. Importantly, current data suggest that every patient who has had a gastric band removed, for any reason, will have weight regain back to or near their baseline level [45].

**Cost effectiveness of gastric banding**

The long term benefits of bariatric surgery for weight loss, co-morbidity resolution, lower cancer rates and improved mortality could be expected to lead to a health care cost saving but results are not that clear cut. While a systematic review in 2011 [46] suggested that weight-loss surgery was cost effective compared to non-surgical treatment, a 2014 study on insurance provider claim costs after bariatric surgery showed no reduction in actual costs of claims [47]. However, the measure used for comparison is important, and many studies in different countries have shown that the incremental cost effectiveness ratio per quality- or disability- adjusted life year varied from about £1,300 for gastric banding in Australia [48, to £4,000 - £4,500 for all laparoscopic weight loss surgery in the USA [49]. Interestingly, while

all these values are far below the cost-effectiveness threshold for the respective countries (and also for the UK), gastric banding may be the most cost saving treatment at a population level.

An analysis of data using a decision analytic model from the Scandinavian Registry, covering cardiovascular disease, diabetes and surgical complications, showed that bariatric surgery led to a cost saving when compared to non-surgical management of weight loss [50]. When the risk reduction of vascular events, reduction in diabetes and gain in quality adjusted life years were extrapolated for the whole cohort, operated on in one year, the lifetime gains were over 32,000 quality-adjusted person years, and a net saving of £43 million in health care costs.

The very low volume of bariatric surgery in the UK represents much less than 1 in 100 of the 5%, or 2.1 million people [51], who meet UK NICE guidelines for surgery. In other developed countries there are large differences in the prevalence of co-morbidities and socio-demographic status between surgery-eligible patients and bariatric surgery recipients [52]. Clearly no healthcare system could provide capacity to treat such a potentially great demand, but the overwhelming evidence is that surgery is clinically beneficial, and cost effective. Thus, the RCT evidence in the NICE Guidance showed clear improvement in every aspect of diabetes control and on this basis increasing availability of bariatric surgery should be a clear aspiration for healthcare services. Alternatively, the availability of government funded weight-loss surgery could be limited to the procedure with the lowest costs and shortest hospitalization requirements, to maximise the opportunity for delivery of a service. This strategy would favour gastric banding, although this does not take into account the longer term follow up costs, and the provision of services required, for gastric band adjustments in the community or specialised clinics.

## CONCLUSIONS

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Obesity, with nearly 30 million sufferers in the UK currently, and its treatment, is a huge problem for the NHS. Current evidence shows that only bariatric surgery gives clinically meaningful weight loss, although there is a paucity of high quality long-term evidence. Despite its effectiveness less than 5,000 bariatric procedures were performed in the NHS in 2013 [12, 13], thus the NHS provides the life-saving and quality of life improving benefit of surgery to less than 1 in 4,000 of the severely obese population in the UK. There would need to be a huge turn-around in obesity service provision for any significant population health gains to be achieved. The nature of which operation would best deliver these gains is still unknown, more high quality studies are needed, but with the lowest reported incremental cost effectiveness ratio per quality adjusted life year, gastric banding may provide the NHS with the most punch for its pound. However, unlike Australia, the UK NHS is not currently well set up to look after large number of gastric band patients and adjust the bands regularly in hospital (outside of clinical trials) or in the community. This issue of gastric band follow-up is probably the reason why many patients and surgeons do not choose gastric banding over other operations in the NHS. In addition, many bands are removed by surgeons in non-specialist centres if the patient develops a problem such as slippage, rather than having it rectified. It would also require a shift in the moral perspective of delivering treatments effective at improving quality of life to those with a significant need [53]. Still a major unknown remains the costs associated with the management of the long-term complications of bariatric surgery, including that of modern gastric banding, which appear comparable for all techniques.



## References

1. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384(9945):766-81.
2. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes research and clinical practice*. 2014;103(2):137-49.
3. De Pergola G, Silvestris F. Obesity as a major risk factor for cancer. *Journal of obesity*. 2013;2013:291546.
4. Herrigel DJ, Moss RA. Diabetes mellitus as a novel risk factor for gastrointestinal malignancies. *Postgraduate medicine*. 2014;126(6):106-18.
5. Korhonen PE, Seppälä T, Järvenpää S, Kautiainen H. Body mass index and health-related quality of life in apparently healthy individuals. *Qual Life Res*. 2014 Feb;23(1):67-74.
6. Adams TD, Davidson LE, Litwin SE, Kolotkin RL, LaMonte MJ, Pendleton RC et al. Health benefits of gastric bypass surgery after 6 years. *Jama*. 2012;308(11):1122-31.
7. Mehta T, Fontaine KR, Keith SW, Bangalore SS, de los Campos G, Bartolucci A et al. Obesity and mortality: are the risks declining? Evidence from multiple prospective studies in the United States. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2014;15(8):619-29.
8. Colquitt JL, Pickett K, Loveman E, Frampton GK. Surgery for weight loss in adults. *The Cochrane database of systematic reviews*. 2014;8:Cd003641.
9. Gloy VL, Briel M, Bhatt DL, Kashyap SR, Schauer PR, Mingrone G et al. Bariatric surgery versus non-surgical treatment for obesity: a systematic review and meta-analysis of randomised controlled trials. *BMJ (Clinical research ed)*. 2013;347:f5934.
10. National Clinical Guideline CG189. National Institute for Health and Clinical Excellence: Guidance. Obesity: Identification, Assessment and Management of Overweight and Obesity in Children, Young People and Adults: Partial Update of CG43. London: National Institute for Health and Care Excellence (UK) Copyright (c) National Clinical Guideline Centre, 2014.
11. Buchwald H, Oien DM. Metabolic/bariatric surgery worldwide 2011. *Obesity surgery*. 2013;23(4):427-36.
12. Welbourn R, Small P, Finlay I, Sareela A, Somers S, Mahawar K et al. The Second National Bariatric Surgery Registry Report to March 2013. Dendrite Clinical Systems; Henley UK. ISBN 978-0-9568154-8-4 (2014).
13. Welbourn R, Fiennes A, Kinsman R, Walton P. The UK National Bariatric Surgery Registry. First Report to 2010. Dendrite Clinical Systems, Henley UK. ISBN 1-903968-27-5 (2011).



14. BOMSS. Consultant Outcomes 2013. 2013. <http://www.bomss.org.uk/surgeon-data/>.

15. Rogers CA, Welbourn R, Byrne J, Donovan JL, Reeves BC, Wordsworth S et al. The By-Band study: gastric bypass or adjustable gastric band surgery to treat morbid obesity: study protocol for a multi-centre randomised controlled trial with an internal pilot phase. *Trials*. 2014;15:53.

16. Burton PR, Yap K, Brown WA, Laurie C, O'Donnell M, Hebbard G, Kalff V, O'Brien PE. Changes in satiety, supra- and infraband transit, and gastric emptying following laparoscopic adjustable gastric banding: a prospective follow-up study. *Obes Surg*. 2011 Feb;21(2):217-23.

17. Moore M, Wainwright P, Hopkins JC. Practice pointer: Primary care management of patients after weight loss surgery. *BMJ* in press

18. Brown W, Korin A, Burton P, O'Brien PE. Laparoscopic adjustable gastric banding. *Aust Fam Physician*. 2009 Dec;38(12):972-6.

19. Carlin AM, Zeni TM, English WJ, Hawasli AA, Genaw JA, Krause KR et al. The comparative effectiveness of sleeve gastrectomy, gastric bypass, and adjustable gastric banding procedures for the treatment of morbid obesity. *Annals of surgery*. 2013;257(5):791-7.

20. Peterli R, Borbely Y, Kern B, Gass M, Peters T, Thurnheer M et al. Early results of the Swiss Multicentre Bypass or Sleeve Study (SM-BOSS): a prospective randomized trial comparing laparoscopic sleeve gastrectomy and Roux-en-Y gastric bypass. *Annals of surgery*. 2013;258(5):690-4

21. Higa K, Ho T, Tercero F, Yunus T, Boone KB. Laparoscopic Roux-en-Y gastric bypass: 10-year follow-up. *Surgery for obesity and related diseases*. 2011;7(4):516-25.

22. Sjostrom L, Narbro K, Sjostrom CD, Karason K, Larsson B, Wedel H et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *The New England journal of medicine*. 2007;357(8):741-52.

23. Sjostrom L. Review of the key results from the Swedish Obese Subjects (SOS) trial - a prospective controlled intervention study of bariatric surgery. *Journal of internal medicine*. 2013;273(3):219-34.

24. Angrisani L, Cutolo PP, Formisano G, Nosso G, Vitolo G. Laparoscopic adjustable gastric banding versus Roux-en-Y gastric bypass: 10-year results of a prospective, randomized trial. *Surgery for obesity and related diseases*. 2013;9(3):405-13.

25. Nguyen NT, Slone JA, Nguyen XM, Hartman JS, Hoyt DB. A prospective randomized trial of laparoscopic gastric bypass versus laparoscopic adjustable gastric banding for the treatment of morbid obesity: outcomes, quality of life, and costs. *Ann Surg*. 2009 Oct;250(4):631-41.

26. Puzziferri N, Roshek TB 3rd, Mayo HG, Gallagher R, Belle SH, Livingston EH. Long-term follow-up after bariatric surgery: a systematic review. *JAMA*. 2014 Sep 3;312(9):934-42.

27. Chang S, Stoll CT, Song J, Varela J, Eagon CJ, Colditz GA. The effectiveness and risks of bariatric surgery: An updated systematic review and meta-analysis, 2003-2012. *JAMA Surgery*. 2014;149(3):275-87.

28. O'Brien PE, MacDonald L, Anderson M, Brennan L, Brown WA. Long-term outcomes after bariatric surgery: fifteen-year follow-up of adjustable gastric banding and a systematic review of the bariatric surgical literature. *Annals of surgery*. 2013;257(1):87-94.
29. Courcoulas AP, Christian NJ, Belle SH, Berk PD, Flum DR, Garcia L et al. Weight change and health outcomes at 3 years after bariatric surgery among individuals with severe obesity. *Jama*. 2013;310(22):2416-25.
30. Buchwald H, Estok R, Fahrbach K, Banel D, Jensen MD, Pories WJ et al. Weight and type 2 diabetes after bariatric surgery: systematic review and meta-analysis. *The American journal of medicine*. 2009;122(3):248-56.e5.
31. Yu J, Zhou X, Li L, Li S, Tan J, Li Y et al. The long-term effects of bariatric surgery for type 2 diabetes: systematic review and meta-analysis of randomized and non-randomized evidence. *Obesity surgery*. 2015;25(1):143-58.
32. Dixon JB, Murphy DK, Segel JE, Finkelstein EA. Impact of laparoscopic adjustable gastric banding on type 2 diabetes. *Obes Rev*. 2012 Jan;13(1):57-67.
33. Dixon JB, O'Brien PE, Playfair J, Chapman L, Schachter LM, Skinner S, Proietto J, Bailey M, Anderson M. Adjustable gastric banding and conventional therapy for type 2 diabetes: a randomized controlled trial. *JAMA*. 2008 Jan 23;299(3):316-23. doi: 10.1001/jama.299.3.316.
34. Edelman S, Ng-Mak DS, Fusco M, Ashton D, Okerson T, Liu Q, Jin J, Dixon JB. Control of type 2 diabetes after 1 year of laparoscopic adjustable gastric banding in the helping evaluate reduction in obesity (HERO) study. *Diabetes Obes Metab*. 2014 Oct;16(10):1009-15
35. Wentworth JM, Playfair J, Laurie C, Ritchie ME, Brown WA, Burton P, Shaw JE, O'Brien PE. Multidisciplinary diabetes care with and without bariatric surgery in overweight people: a randomised controlled trial. *Lancet Diabetes Endocrinol*. 2014 Jul;2(7):545-52.
36. Carlsson LM, Peltonen M, Ahlin S, Anveden A, Bouchard C, Carlsson B et al. Bariatric surgery and prevention of type 2 diabetes in Swedish obese subjects. *The New England journal of medicine*. 2012;367(8):695-704.
37. Himpens J, Cadière GB, Bazi M, Vouche M, Cadière B, Dapri G. Long-term outcomes of laparoscopic adjustable gastric banding. *Arch Surg*. 2011 Jul;146(7):802-7.
38. Aminian A, Brethauer SA, Kirwan JP, Kashyap SR, Burguera B, Schauer PR. How safe is metabolic/diabetes surgery? *Diabetes, obesity & metabolism*. 2015;17(2):198-201.
39. Hopkins JC, Howes N, Chalmers K, Savovic J, Whale K, Coulman KD et al. Outcome reporting in bariatric surgery: an in-depth analysis to inform the development of a core outcome set, the BARIACT Study. *Obesity reviews*. 2015;16(1):88-106.
40. Sudan R, Nguyen NT, Hutter MM, Brethauer SA, Ponce J, Morton JM. Morbidity, mortality, and weight loss outcomes after reoperative bariatric surgery in the USA. *Journal of gastrointestinal surgery*. 2015;19(1):171-8; discussion 8-9.

41. Karmali S, Brar B, Shi X, Sharma AM, de Gara C, Birch DW. Weight recidivism post-bariatric surgery: a systematic review. *Obesity surgery*. 2013;23(11):1922-33.

42. te Riele WW, Boerma D, Wiezer MJ, Borel Rinkes IH, van Ramshorst B. Long-term results of laparoscopic adjustable gastric banding in patients lost to follow-up. *Br J Surg*. 2010 Oct;97(10):1535-40.

43. Hochberg LS, Murphy KD, O'Brien PE, Brennan L. Laparoscopic Adjustable Gastric Banding (LAGB) Aftercare Attendance and Attrition. *Obes Surg*. 2015 Sep;25(9):1693-702.

44. Sysko R, Hildebrandt TB, Kaplan S, Brewer SK, Zitsman JL, Devlin MJ. Predictors and correlates of follow-up visit adherence among adolescents receiving laparoscopic adjustable gastric banding. *Surg Obes Relat Dis*. 2014 Sep-Oct;10(5):914-20.

45. Aarts EO, Dogan K, Koehestanie P, Janssen IM, Berends FJ. What happens after gastric band removal without additional bariatric surgery? *Surg Obes Relat Dis*. 2014 Nov-Dec;10(6):1092-6.

46. Padwal R, Klarenbach S, Wiebe N, Hazel M, Birch D, Karmali S et al. Bariatric surgery: a systematic review of the clinical and economic evidence. *Journal of general internal medicine*. 2011;26(10):1183-94.

47. Weiner JP, Goodwin SM, Chang HY, Bolen SD, Richards TM, Johns RA et al. Impact of bariatric surgery on health care costs of obese persons: a 6-year follow-up of surgical and comparison cohorts using health plan data. *JAMA Surg*. 2013;148(6):555-62.

48. Lee YY, Veerman JL, Barendregt JJ. The cost-effectiveness of laparoscopic adjustable gastric banding in the morbidly obese adult population of Australia. *PloS one*. 2013;8(5):e64965.

49. Wang BC, Wong ES, Alfonso-Cristancho R, He H, Flum DR, Arterburn DE et al. Cost-effectiveness of bariatric surgical procedures for the treatment of severe obesity. *The European journal of health economics:HEPAC* 2014;15(3):253-63.

50. Borisenko O, Adam D, Funch-Jensen P, Ahmed AR, Zhang R, Colpan Z et al. Bariatric Surgery can Lead to Net Cost Savings to Health Care Systems: Results from a Comprehensive European Decision Analytic Model. *Obesity surgery*. 2015 Sep;25(9):1559-68.

51. Ahmad A, Lavery AA, Aasheim E, Majeed A, Millett C, Saxena S. Eligibility for bariatric surgery among adults in England: analysis of a national cross-sectional survey. *JRSM Open*. 2014 Jan 7;5(1):2042533313512479. doi: 10.1177/2042533313512479. eCollection 2014.

52. Padwal RS, Chang HJ, Klarenbach S, Sharma AM, Majumdar SR. Characteristics of the population eligible for and receiving publicly funded bariatric surgery in Canada. *International journal for equity in health*. 2012;11:54.

53. Persson K. Why Bariatric surgery should be given high priority: an argument from law and morality. *Health care analysis:HCA: journal of health philosophy and policy*. 2014;22(4):305-24.



Dear Mr Richard Welbourn

We hereby grant you permission to reproduce the material detailed below in **print** and **electronic** format at no charge subject to the following conditions:

1. If any part of the material to be used (for example, figures) has appeared in our publication with credit or acknowledgement to another source, permission must also be sought from that source. If such permission is not obtained then that material may not be included in your publication/copies.

2. Suitable acknowledgement to the source must be made, either as a footnote or in a reference list at the end of your publication, as follows:

“This article was published in Publication title, Vol number, Author(s), Title of article, Page Nos, Copyright Elsevier (or appropriate Society name) (Year).”

3. This permission is granted for non-exclusive world rights in all languages.

4. Reproduction of this material is granted for the purpose for which permission is hereby given.

Yours sincerely



Jennifer Jones

Permissions Specialist

Elsevier Limited, a company registered in England and Wales with company number 1982084, whose registered office is The Boulevard, Langford Lane, Kidlington, Oxford, OX5 1GB, United Kingdom.